

APPENDICES
FOR
UNITED STATES LETTERS PATENT

TITLE: **OVERCOMING DAPA AMINOTRANSFERASE BOTTLENECKS
IN BIOTIN VITAMERS BIOSYNTHESIS**

APPLICANT: **SCOTT W. VAN ARSDELL, R. ROGERS YOCUM, JOHN B.
PERKINS, and JANICE G. PERO**

2 PAGES OF APPENDICES

"EXPRESS MAIL." Mailing Label Number EM526114158105

Date of Deposit July 14, 1997

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

USA G. Gray
USA G. Gray

Appendix I. Medium composition for biotin and vitamins production in bench scale fermentors.

Medium Component	Batch	Concentration	Feed
Glucose	15.0 g/liter		750 g/liter
Veal Infusion Broth ¹	25.0 g/liter		---
Yeast Extract ¹	5.0 g/liter		---
Sodium Glutamate	5.0 g/liter		---
KH ₂ PO ₄	7.5 g/liter		13.7 g/liter
MgCl ₂ ·6H ₂ O	1.0 g/liter		1.5 g/liter
(NH ₄) ₂ SO ₄	2.0 g/liter		---
MAZU DF-37C	2.5 g/liter		---
CaCl ₂ ·2H ₂ O	1.0 g/liter		---
CuSO ₄ ·5H ₂ O	0.4 mg/liter		4.0 mg/liter
ZnSO ₄ ·7H ₂ O	0.5 mg/liter		5.0 mg/liter
MnSO ₄ ·H ₂ O	25.0 mg/liter		35.0 mg/liter
FeSO ₄ ·6H ₂ O	1.0 mg/liter		10.0 mg/liter
Sodium Molybdate-2H ₂ O	0.2 mg/liter		2.0 mg/liter
FeSO ₄ ·7H ₂ O	50.0 mg/liter		100.0 mg/liter
Sodium Citrate-2H ₂ O	50.0 mg/liter		100.0 mg/liter

¹ In Amberex Medium the Veal Infusion Broth and Yeast Extract are replaced with 10 g/l Amberex 695.

Appendix II. Protocol of avidin-HABA [2-(4-hydroxyphenylazo) benzoic acid] displacement assay for biotin and dethiobiotin.

Reagents and Solutions:

Buffer: 0.1 M NaPO₄, pH 7.0.
Avidin: From Sigma (Cat # A-9275). Dissolved at 5 mg/ml in Buffer.
HABA: From Aldrich (Cat # 14,803-2). Dissolved at 0.375 M in water + 1 eq. NaOH.

Prepare Mix:

	20 samples	50 samples
Avidin	1 ml	2.5 ml
HABA	0.08 ml	0.2 ml
Buffer	38.9 ml	97.3 ml

Assay:

Zero spectrophotometer;

Add 2 ml of Buffer to disposable 5 ml cuvette; record OD₅₀₀.

To read sample:

Place disposable 5 ml cuvette in spectrophotometer.

Add 2 ml of Mix; stir; record OD₅₀₀.

Add sample in 0.1 ml volume; stir; record OD₅₀₀.

Standards:

Use 0.1 ml DTB at 2 mg/ml to 14 mg/ml as samples.

Use 0.1 ml Buffer as "zero" point.

Calculations:

Calculate ΔOD_{500} minus ΔOD_{500} .

Plot standards and use curve to determine HABA vitamers from samples.

- Notes:**
1. Useful range is 2 to 14 mg/l of biotin + dethiobiotin.
 2. Add mix to cuvette, read OD₅₀₀, and then add sample and mix without removing cuvette from the spectrophotometer.
 3. Best results are obtained when a constant volume is used with a set of samples and standards. Use Buffer to bring all samples to the same volume.